

STA 3000, Problem Set 4. Due in class December 5.

1. Chapter 3, exercise 12.
2. Chapter 3, exercise 14. Hint for (a): Find a parameterized family of decision rules that includes  $\delta(x) = x$ , and also includes a rule that dominates it.
3. Chapter 3, exercise 17.
4. Do the following:
  - (a) Show that if  $\lambda_1 \ll \lambda_2$  and  $\lambda_2 \ll \lambda_1$ , then any decision rule that is  $\lambda_1$ -admissible is also  $\lambda_2$ -admissible.
  - (b) Give an example of a decision rule  $\delta$  and measures  $\lambda_1$  and  $\lambda_2$  for which  $\lambda_1 \ll \lambda_2$  and  $\delta$  is  $\lambda_1$ -admissible, but  $\delta$  is not  $\lambda_2$ -admissible.
  - (c) Give an example of a decision rule  $\delta$  and measures  $\lambda_1$  and  $\lambda_2$  for which  $\lambda_2 \ll \lambda_1$  and  $\delta$  is  $\lambda_1$ -admissible, but  $\delta$  is not  $\lambda_2$ -admissible.
5. Suppose a positive integer  $X$  is observed from a distribution of the following form:

$$P_\theta(X = x) = \begin{cases} 3/4 & \text{if } x = \theta \\ 1/4 & \text{if } x = \theta + 1 \\ 0 & \text{otherwise} \end{cases}$$

where the parameter  $\theta$  is a positive integer (ie,  $\Omega = \{1, 2, 3, \dots\}$ ). We wish to find a decision rule that estimates  $\theta$  with 0–1 loss — ie,

$$L(\theta, \delta(x)) = \begin{cases} 0 & \text{if } \delta(x) = \theta \\ 1 & \text{if } \delta(x) \neq \theta \end{cases}$$

- (1) Find the risk functions for each of the estimators  $\delta_1(x) = x$ ,  $\delta_2(x) = x + 1$ , and  $\delta_3(x) = x - 1$ .
- (2) Find a Bayes rule for each of the following prior distributions:
  - (a)  $P(\Theta = \theta) = 2^{-\theta}$
  - (b)  $P(\Theta = \theta) = 3 \cdot 4^{-\theta}$
  - (c)  $P(\Theta = \theta) = 2 \cdot 3^{-\theta}$

Determine whether or not each of these rules is admissible.

- (3) Which of the rules  $\delta_1(x) = x$ ,  $\delta_2(x) = x + 1$ , and  $\delta_3(x) = x - 1$  are admissible?
- (4) Find a minimax rule for this decision problem.