

The Bayesian Approach to Machine Learning

- 1) We formulate our knowledge about the situation probabilistically:
 - We define a *model* that expresses qualitative aspects of our knowledge (eg, forms of distributions, independence assumptions). The model will have some unknown *parameters*.
 - We specify a *prior* probability distribution for these unknown parameters that expresses our beliefs about which values are more or less likely, before seeing the data.
- 2) We gather data.
- 3) We compute the *posterior* probability distribution for the parameters, given the observed data.
- 4) We use this posterior distribution to make predictions by averaging over the posterior distribution, or to make decisions so as to minimize posterior expected loss.

Finding the Posterior Distribution

The *posterior distribution* for the model parameters given the observed data is found by combining the prior distribution with the likelihood for the parameters given the data.

This is done using *Bayes' Rule*:

$$P(\text{parameters} \mid \text{data}) = \frac{P(\text{parameters}) P(\text{data} \mid \text{parameters})}{P(\text{data})}$$

The denominator is just the required normalizing constant, and can often be filled in at the end, if necessary. So as a proportionality, we can write

$$P(\text{parameters} \mid \text{data}) \propto P(\text{parameters}) P(\text{data} \mid \text{parameters})$$

which can be written schematically as

$$\text{Posterior} \propto \text{Prior} \times \text{Likelihood}$$

We make predictions by integrating with respect to the posterior:

$$P(\text{new data} \mid \text{data}) = \int_{\text{parameters}} P(\text{new data} \mid \text{parameters}) P(\text{parameters} \mid \text{data})$$

Representing the Prior and Posterior Distributions by Samples

The complex distributions we will often use as priors, or obtain as posteriors, may not be easily represented or understood using formulas.

A very general technique is to represent a distribution by a *sample* of many values drawn randomly from it. We can then:

- *Visualize* the distribution by viewing these sample values, or low-dimensional projections of them.
- Make *Monte Carlo* estimates for probabilities or expectations with respect to the distribution, by taking averages over these sample values.

Obtaining a sample from the prior is often easy. Obtaining a sample from the posterior is usually more difficult — but this is nevertheless the dominant approach to Bayesian computation.

A Simple Example — A Hard Linear Classifier

The problem:

We will be observing pairs (x_i, y_i) , for $i = 1, \dots, N$, where $x_i = (x_{i1}, x_{i2})$ are two input variables, and y_i is a $-1/ +1$ class indicator. We are interested in predicting y from x for a test case.

Our informal beliefs:

We believe that there is a line somewhere in the input space that determines y perfectly — with -1 on one side, $+1$ on the other.

We think that this line could equally well have any orientation, and that it could equally well be positioned anywhere, as long as it is no more than a distance of three from the origin at its closest point.

We need to translate these informal beliefs into a *model* and a *prior*.

Formalizing the Model

Our model can be formalized by saying that

$$P(y_i = y \mid x_i, u, w) = \begin{cases} 1 & \text{if } y u (w^T x_i - 1) > 0 \\ 0 & \text{if } y u (w^T x_i - 1) < 0 \end{cases}$$

where $u \in \{-1, +1\}$ and $w = (w_1, w_2)$ are unknown *parameters* of the model. The value of w determines a line separating the classes, and u says which class is on which side.


This model is rather dogmatic — eg, it says that y is **certain** to be $+1$ if $u = +1$ and $w^T x$ is greater than 1. A more realistic model would replace the probabilities of 0 and 1 above with ϵ and $1 - \epsilon$ to account for possible unusual items, or for misclassified items. ϵ might be another unknown parameter.

Formalizing the Prior

A line is completely determined by giving the point, c , on the line that is closest to the origin.

To formalize our prior belief that the line separating classes could equally well be anywhere, as long as it is no more than a distance of three from the origin, we decide to use a uniform distribution for c over the circle with radius 3.

Given c , we can compute $w = c/\|c\|^2$, which makes $w^T x = 1$ for points on the line. (Here, $\|c\|^2$ is the squared norm, $c_1^2 + c_2^2$.)




Here's an example:

We also say that w is equally likely to be +1 or -1, independently of w .

Looking at the Prior Distribution


We can check this prior distribution by looking at many lines sampled from it:



Something's wrong here. We meant for the lines to be uniformly distributed, but we see a sparse region near the origin.

Why This Prior Distribution is Wrong

Our first attempt at formalizing our prior beliefs didn't work. We can see why if we think about it.




Imagine moving a line that's within five degrees of vertical from left to right:

To stay within five degrees of vertical, the closest point to the origin has to be within the wedge shown. This becomes less and less likely as the origin is approached. We don't get the same probability of a near-vertical line for all horizontal positions.

Fixing the Prior Distribution


We can fix the prior by letting the closest point on the line to the origin be $c = ru$, with r uniformly distributed over $(0, 3)$ and u uniformly distributed over the unit circle.

Now a sample drawn from the prior looks the way we want it to:



Some Data Points

Now that we have defined our model and prior, let's get some data:



The black points are in class +1, the white points in class -1.

Posterior Distribution for the Hard Linear Classifier

For the hard linear classifier, the likelihood is either 0 or 1:


$$\begin{aligned} P(y_1, \dots, y_N \mid x_1, \dots, x_N, u, w) &= \prod_{i=1}^N P(y_i \mid x_i, u, w) \\ &= \begin{cases} 1 & \text{if } y_i u (w^T x_i - 1) > 0, \text{ for } i = 1, \dots, N \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

The posterior distribution for u and w is therefore the same as their prior distribution, except that parameter values incompatible with the data are eliminated.

After renormalizing so that posterior probabilities integrate to one, the parameter values compatible with the data will have higher probability than they did in the prior.

Obtaining a Sample from the Posterior Distribution

To obtain a sample of values from the posterior, we can sample w values from the prior, but retain only those that are compatible with the data (for some u).
Here's what we get using a sample of size 200:



The eight bold lines are a random sample from the posterior distribution.

Making a Prediction for a Test Case

The Bayesian predictive probability that in a test case with inputs x_* , the class, y_* , will be +1 is found by integrating/summing over the parameters w and u :

$$\begin{aligned} P(y_* = +1 \mid x_*, (x_1, y_1), \dots, (x_N, y_N)) \\ = \int \sum_{u=\pm 1} P(y_* = +1 \mid x_*, u, w) P(u, w \mid x_1, y_1), \dots, (x_N, y_N) dw \end{aligned}$$


Using a sample of K values from the posterior, $(u^{(1)}, w^{(1)}), \dots, (u^{(K)}, w^{(K)})$, we can approximate this as follows:

$$P(y_* = +1 \mid x_*, (x_1, y_1), \dots, (x_N, y_N)) \approx \frac{1}{K} \sum_{j=1}^K P(y_* = +1 \mid x_*, u^{(j)}, w^{(j)})$$

For this model, $P(y_* = +1 \mid x_*, u^{(j)}, w^{(j)})$ is either 0 or 1, depending on the sign of $u^{(j)} (w^{(j)})^T x_* - 1)$. The average above is just the fraction of lines drawn from the posterior that would put the test point in class +1.

A Plot of the Predictive Probabilities

Here is a contour plot over the input space of the approximate predictive probability of class +1, based on a sample of size 10000 from the prior, which resulted in a sample of size 450 from the posterior:



The contour lines go from 0 on the left to 1 on the right, in steps of 0.1.

Final Thoughts On This Example

- We see that correctly translating informal knowledge into a prior distribution isn't always trivial.
- However, a prior can be *tested*, by checking how well it corresponds to our prior beliefs. Prior distributions are **not** “arbitrary”.
- More elaborate priors might sometimes be appropriate. For example, we might use a prior that favoured lines that are almost horizontal or vertical, if we believe that probably one of the two inputs is mostly irrelevant.
- For a data set with seven points, only about 4.5% of the parameter values we drew from the prior made it into the posterior sample. This technique isn't going to work for realistic problems. We need better ways of sampling from the posterior distribution.